

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. §371**

SCH 1851

U.S. APPLICATION NO. (If known, see 37 CFR §1.5)

10/018429

INTERNATIONAL APPLICATION NO

PCT/EP00/05969

INTERNATIONAL FILING DATE

26 JUNE 2000

PRIORITY DATE CLAIMED

24 JUNE 1999

TITLE OF INVENTION

11β LONG-CHAIN SUBSTITUTED ESTRATRIENES, METHOD FOR THEIR PRODUCTION, PHARMACEUTICAL PREPARATIONS CONTAINING SAID 11β LONG-CHAIN SUBSTITUTED ESTATRIENES, AND THEIR USE FOR PRODUCING MEDICAMENTS.

APPLICANT(S) FOR DO/EO/US


BOHLMANN, Rolf, et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. §371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. §371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. §371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. §371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. §371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 C.F.R. §§1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. §§3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

U.S. APPLICATION NO. (if known, see 37 CFR §1.5) <div style="font-size: 2em; font-weight: bold; margin-top: 5px;">10/018429</div>		INTERNATIONAL APPLICATION NO. PCT/EP00/05969		ATTORNEY'S DOCKET NUMBER SCH 1851	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR §1.492 (a) (1) - (5)): Search Report has been prepared by the EPO or JPO..... \$890.00 International preliminary examination fee paid to USPTO (37 CFR §1.482)..... \$710.00 No international preliminary examination fee paid to USPTO (37 CFR §1.482) but international search fee paid to USPTO (37 CFR §1.445(a)(2))..... \$740.00 Neither international preliminary examination fee (37 CFR §1.482) nor international search fee (37 CFR §1.445(a)(2)) paid to USPTO..... \$1040.00 International preliminary examination fee paid to USPTO (37 CFR §1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... \$100.00				CALCULATIONS PTO USE ONLY	
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$890.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 C.F.R. §1.492(e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	8 - 20 =	0	x \$ 18.00	\$0.00	
Independent claims	2 - 3 =	0	x \$ 84.00	\$0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)				+ \$ 280.00	
TOTAL OF ABOVE CALCULATIONS =				\$890.00	
Reduction of 1/2 for filing by small entity, if applicable. A Verified Small Entity Statement must also be					
SUBTOTAL =				\$890.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 C.F.R. §1.492(f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
TOTAL NATIONAL FEE =				\$890.00	
Fee for recording the enclosed assignment (37 C.F.R. §1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. §§3.28, 3.31). \$40.00 per property.					
TOTAL FEES ENCLOSED =				\$890.00	
				Amount to be refunded:	
				charged:	
<p>a. <input checked="" type="checkbox"/> A check in the amount of <u>\$890.00</u> to cover the above fees is enclosed.</p> <p>b. <input type="checkbox"/> Please charge my Deposit Account No. <u>13-3402</u> in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed.</p> <p>c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-3402</u>. A duplicate copy of this sheet is enclosed.</p> <p>NOTE: Where an appropriate time limit under 37 C.F.R. §§1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. §1.137(a) or (b)) must be filed and granted to restore the application to pending status.</p> <p>SEND ALL CORRESPONDENCE TO: Customer Number 23,599</p> <div style="display: flex; justify-content: space-between; align-items: flex-end; margin-top: 20px;"> <div style="text-align: center;">  <div style="font-size: 1.5em; font-weight: bold;">23599</div> <div style="font-size: 0.8em;">PATENT TRADEMARK OFFICE</div> </div> <div style="text-align: right;"> <div style="font-size: 1.2em; font-family: cursive;">Anthony J. Zelano</div> <div style="font-size: 0.8em; margin-top: 5px;">SIGNATURE</div> <div style="border-bottom: 1px solid black; width: 150px; margin: 5px 0;"></div> <div style="font-size: 0.8em; margin-top: 5px;">NAME</div> <div style="border-bottom: 1px solid black; width: 100px; margin: 5px 0;"></div> <div style="font-size: 0.8em; margin-top: 5px;">REGISTRATION NUMBER</div> </div> </div>					

Filed: 20 DECEMBER 2001
AJZ:kmo

Anthony J. Zelano
NAME
27,969
REGISTRATION NUMBER

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
4. Januar 2001 (04.01.2001)

PCT


(10) Internationale Veröffentlichungsnummer
WO 01/00652 A2

- (51) Internationale Patentklassifikation⁷: C07J (72) Erfinder; und
(75) Erfinder/Anmelder (nur für US): BOHLMANN, Rolf [DE/DE]; Kühler Weg 6A, D-14055 Berlin (DE). HEINRICH, Nikolaus [DE/DE]; Sponholzstrasse 8, D-12159 Berlin (DE). KROLL, Jorg [DE/DE]; Horst-Kohl-Strasse 3, D-12157 Berlin (DE). SAUER, Gerhard [DE/DE]; Königsbacher Zeile 41a, D-13465 Berlin (DE).
- (21) Internationales Aktenzeichen: PCT/EP00/05969
- (22) Internationales Anmeldedatum: 26. Juni 2000 (26.06.2000)
- (25) Einreichungssprache: Deutsch (81) Bestimmungsstaaten (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (26) Veröffentlichungssprache: Deutsch
- (30) Angaben zur Priorität: 199 29 715.0 24. Juni 1999 (24.06.1999) DE (84) Bestimmungsstaaten (regional): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
- (71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): SCHERING AKTIENGESELLSCHAFT [DE/DE]; Müllerstrasse 178, D-133353 Berlin (DE).

[Fortsetzung auf der nächsten Seite]

(54) Title: 11 β LONG-CHAIN SUBSTITUTED ESTRADIENES, METHOD FOR THEIR PRODUCTION, PHARMACEUTICAL PREPARATIONS CONTAINING SAID 11 β LONG-CHAIN SUBSTITUTED ESTRADIENES, AND THEIR USE FOR PRODUCING MEDICAMENTS

(54) Bezeichnung: 11 β -LANGKETTIG-SUBSTITUIERTE ESTRADIENE, VERFAHREN ZUR HERSTELLUNG, PHARMAZEUTISCHE PRÄPARATE, DIE DIESE 11 β -LANGKETTIG-SUBSTITUIERTEN ESTRADIENE ENTHALTEN, SOWIE DAREN VERWENDUNG ZUR HERSTELLUNG VON ARZNEIMITTELEN



(57) Abstract: The invention relates to novel 11 β long-chain substituted estradienes of general formula (I), wherein R¹¹ is a long-chain residue with one nitrogen and optionally one sulfur atom, that can additionally be terminally functionalized with a perfluoroalkyl group or an optionally substituted aryl group. The inventive compounds are characterized by antiestrogenic or tissue-selective estrogenic properties and are useful in the production of medicaments.

(57) Zusammenfassung: Die vorliegende Erfindung beschreibt die neuen 11 β -langkettig-substituierten Estradiene der allgemeinen Formel (I), worin R¹¹ ein langkettiger, ein Stickstoff- sowie gegebenenfalls ein Schwefelatom aufweisender Rest ist, der außerdem endständig mit einer Perfluoralkylgruppe oder einem gegebenenfalls substituierten Arylrest funktionalisiert sein kann. Die Verbindungen verfügen über antiestrogene oder gewebe selektive estrogenische Eigenschaften und sind zur Herstellung von Arzneimitteln geeignet.

WO 01/00652 A2


10/018429

WO 01/00652

PCT/EP00/05969

11 β -Long-Chain-Substituted Estratrienes, Process for the
Production, Pharmaceutical Preparations that Contain these 11 β -
Long-Chain-Substituted Estratrienes, as well as their Use for the
Production of Pharmaceutical Agents.

This invention relates to 11 β -long-chain-substituted estratrienes of general formula I



(I)

in which

R³ means a hydrogen atom, a hydrocarbon radical with up to 8 carbon atoms or a radical of partial formula R^{3'}-C(O)-, in which R^{3'} means a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms or a phenyl radical,

R¹¹ means a radical of formula -A-B-Z-R²⁰,
in which

A stands for a direct bond, and

- B stands for a straight-chain or branched-chain alkylene, alkenylene or alkynylene group with 4, 5 or 6 carbon atoms, or
- A stands for a phenylene radical, and
- B stands for a methylene, ethylene, propylene or trimethylene group, or
- A stands for a phenylenoxy radical, whereby the latter is bonded via a carbon atom to the 11-carbon atom of the steroid, and
- B stands for an ethylene group, and
- Z stands for $-NR^{21}-$ and R^{21} stands for a C_1-C_3 alkyl group,

whereby R^{20} means

a hydrogen atom,
 a straight-chain or branched-chain alkyl, alkenyl or alkynyl group with up to 10 carbon atoms, whereby if A is a direct bond, R^{20} and R^{21} do not both simultaneously mean methyl, however, and, if A is a phenylenoxy radical, R^{20} and R^{21} do not both simultaneously mean methyl or ethyl in each case, and if A is a phenylenoxy radical and B means an ethylene group, OR^{17b} should not be a hydroxy group and R^{17a} should not be a C_{1-4} alkyl group, and R^3 should not be a hydrogen atom,

or one of groupings

10048429-050702

$-D-C_nF_{2n+1}$, whereby D is a straight-chain or branched-chain alkylene, alkenylene or alkynylene group with up to 8 carbon atoms and n is an integer from 1 to 8,

D-aryl, whereby D has the already indicated meaning, and aryl stands for a phenyl, 1- or 2-naphthyl radical or a heteroaryl radical that is optionally substituted in one or two places,

$-L-CH=CF-C_pF_{2p+1}$, whereby L is a straight-chain or branched-chain alkylene, alkenylene or alkynylene group with up to 7 carbon atoms and p is an integer from 1 to 7,

whereby in the three cases above in D or L, a methylene group can be replaced by a sulfur atom, a sulfone group or a sulfoxide group,

$-D-O-(CH_2)_q$ -aryl, whereby D and aryl have the already indicated meanings, and q is 0, 1, 2 or 3,

$-D-O-(CH_2)_r-C_nF_{2n+1}$, whereby D and n have the already indicated meanings, and r stands for an integer from 1 to 5,

whereby in addition in all relevant cases above, R^{21} together with D with the inclusion of the nitrogen atom can

then form a pyrrolidine ring that is substituted in 2- or 3-position,

or

if A is a direct bond or a phenylene radical, R^{20} and R^{21} with the nitrogen atom to which they are bonded form a saturated or unsaturated heterocyclic compound with 5 or 6 chain links, which optionally contains one or two additional heteroatoms, selected from nitrogen, oxygen and sulfur, and optionally is substituted,

whereby if A is a phenylene radical and B is a trimethylene radical, R^{21} and R^{20} do not form a methyl or ethyl group, or, together with the nitrogen atom to which they are bonded, do not form a pyrrolidine or piperidine ring,

and

R^{17a} in α - or β -position means a hydrogen atom, a C_{1-5} alkyl, a C_{2-5} alkenyl or a C_{2-5} alkinyl group or a trifluoromethyl group, or together with the radical OR^{17b} means a keto-oxygen atom, and

R^{17b} means a hydrogen atom or a radical of partial formula $R^{17'}-C(O)-$, in which $R^{17'}$ means a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms.

As R^3 , the substituted estratrienes according to the invention preferably have a hydrogen atom. The hydroxy group, however, can also be etherified with a straight-chain or

202050-6248T001

branched-chain, saturated or unsaturated hydrocarbon radical with up to 8 carbon atoms, such as, e.g., a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl, heptyl, hexyl or octyl radical or esterified with an acyl radical $R^{3'}-C(O)-$, in which $R^{3'}$ is a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms or a phenyl radical.

A hydrogen atom or a radical of partial formula $R^{17'}-C(O)-$ can stand for substituents R^{17b} , in which $R^{17'}$ is a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms. A hydrogen atom is preferred for R^{17b} . The hydrocarbon radical can have the meaning of, for example, a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl, heptyl, hexyl or octyl radical. In addition, the substituent $-OR^{17b}$ can be in α - or β -position. The β -position is preferred.

R^{17b} can mean a hydrogen atom, a straight-chain or branched C_{1-5} alkyl radical, such as, for example, a methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl radical, a straight-chain or branched C_{2-5} alkenyl radical, such as, for example, an ethenyl, 1-propenyl, 2-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-ethyl-ethenyl, 2-ethylethenyl, 1-methyl(1-propenyl), 1-methyl(2-propenyl) radical, or a straight-chain or branched C_{2-5} alkynyl radical, such as, for example, an ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-butylnyl, 3-butylnyl, 3-methyl(1-butylnyl)-, 1-methyl(3-butylnyl) radical and a trifluoromethyl radical.

10018429-050702
202050-6484001

R^{17b} preferably means a hydrogen atom, a C_{2-3} alkenyl radical, a C_{2-3} alkynyl radical or a trifluoromethyl group.

R^{17a} especially preferably means a hydrogen atom, a methyl group, an ethenyl radical, an ethynyl radical or a trifluoromethyl group.

In addition, the radical R^{17a} can be in α - or β -position. The α -position is preferred for R^{17a} .

Another meaning for R^{17b} together with OR^{17a} is a keto-oxygen atom. This meaning is to be preferred before any other substitution in 17-position.

In the compounds of general formula I according to the invention, A stands for a direct bond, a phenylene or phenylenoxy radical, whereby the latter is connected via one of its carbon atoms to carbon atom 11 of the steroid skeleton.

An aryl radical that optionally can be substituted is a phenyl, 1- or 2-naphthyl radical in terms of this invention; the phenyl radical is preferred. Unless expressly indicated otherwise, aryl also always includes a heteroaryl radical. Examples of a heteroaryl radical are the 2-, 3- or 4-pyridinyl, the 2- or 3-furyl, the 2- or 3-thienyl, the 2- or 3-pyrrolyl, the 2-, 4- or 5-imidazolyl, the pyrazinyl, the 2-, 4- or 5-pyrimidinyl or the 3- or 4-pyridazinyl radical.


If R^{20} and R^{21} with the nitrogen atom, to which they are bonded, contain a saturated or unsaturated heterocycle with 5 or 6 chain links, which optionally contains one or two additional heteroatoms that are selected from nitrogen, oxygen and sulfur,


this is especially a pyrrolidine, piperidine, morpholine or piperazine ring.

As substituents for the aryl, heteroaryl, aralkyl and heteroarylalkyl radicals, for example, a methyl-, ethyl-, propyl-, trifluoromethyl-, pentafluoroethyl-, trifluoromethylthio-, methoxy-, ethoxy-, nitro-, cyano-, halogen- (fluorine, chlorine, bromine, iodine), hydroxy-, amino-, mono(C_{1-8} alkyl)- or di(C_{1-8} alkyl)amino, whereby both alkyl groups are identical or different, di(aralkyl)amino, whereby both aralkyl groups are identical or different (for aralkyl, see above at R^{20} and R^{31}) or the 1-methoxyacetyl amino radical can be mentioned.

The sulfur atom in the side chain can be present as a single sulfur bridge (sulfide), as sulfone or sulfoxide.

As specific side chains,





can be mentioned.

[Key:]

$-(CH_2)_5-[2-Pyrrolidin-1-yl]-\dots = (CH_2)_5-[2-pyrrolidine-1-yl]-\dots$

p-Phylen-... = p-phenylene-...

This invention relates to the following compounds, i.a.:

11 β -[5-(Methyl(3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl)amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}amino)pentyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfanyl-methyl}pyrrolidin-1-yl]pentyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfinyl-methyl}pyrrolidin-1-yl]pentyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}amino)ethyl]phenyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfinyl]propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

10016429.050702

11 β -{4-[2-(methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}-amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}-amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(8,8,9,9,9-pentafluoro-nonyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-nonyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(9,9,10,10,10-pentafluoro-decyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(8,8,9,9,9-pentafluoro-nonyl)amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(9,9,10,10,10-pentafluoro-decyl)-amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-(methyl-amino)-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-pyrrolidin-1-yl-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4,4,5,5,5-pentafluoro-pentyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4,4,5,5,6,6,7,7,8,8,9,9,9-tridecafluoro-nonyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

202050 62481001

11 β -{5-[(4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11,11-heptadecafluoro-undecyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3,3,4,4,5,5,6,6,6-nonafluoro-hexyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,8-pentafluoro-octyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(7,7,8,8,8-pentafluoro-octyl)-amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,9,9,10,10,10-nonafluoro-decyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,9,9,10,10,11,11,12,12,12,12-tridecafluoro-dodecyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(7,7,8,8,9,9,10,10,11,11,12,12,13,13,14,14,14-heptadecafluoro-tetradecyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,5-hexafluoro-pent-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,6,6,7,7,8,8,8-dodecafluoro-oct-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-hexadecafluoro-dec-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-phenoxy-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

10049429-050703

11 β -{5-[(3-benzyloxy-propyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[N-methyl-N-3-(4,4,5,5,5-pentafluoropentyloxy)-propylamino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -[9-(4,4,5,5,5-pentafluoropentylsulfinyl)-nonyl]-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(2-p-tolyl-ethyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[2-(4-ethoxy-phenyl)-ethyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-phenyl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-pyridin-3-yl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-p-tolyl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[3-(4-chloro-phenyl)-propyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[3-(4-ethoxy-phenyl)-propyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4-methyl-pentyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

In addition to these compounds of general formula I, this invention also relates to their physiologically compatible addition salts with organic and inorganic acids, pharmaceutical preparations that contain these compounds of general formula I

202050-62481001

inclusive of the addition salts, as well as their use for the production of pharmaceutical agents.

Inorganic and organic acids, as are known to one skilled in the art for the formation of physiologically compatible salts, are suitable for the formation of acid addition salts. As addition salts with acids, especially hydrochlorides, hydrobromides, acetates, citrates, oxalates, tartrates and methanesulfonates can be cited.

The compounds of general formula I represent compounds with strong antiestrogenic action and with surprising possible oral uses.

The compounds according to the invention are either pure antiestrogens or so-called partial antagonists, i.e., antiestrogens with partial estrogenic action such as tamoxifen or raloxifen. In contrast to the tamoxifen, their agonistic, estrogenic action is expressed in a tissue-selective manner in the case of partial antagonists of general formula I. In particular, the agonistic action occurs in bone, in the cardiovascular system and in the central nervous system. In particular, no action or only slightly agonistic action occurs in the uterus.

Compounds with antiestrogenic properties, i.e., substances with inhibiting actions compared to estrogens, have already been described extensively.

Estratrienes that carry a β -position substituent in 11-position and that also have, i.a., antiestrogenic action, are

known from, for example, the following patent applications:

WO 98/28324, EP-A 0 850 647, EP-A 0 629 635, WO 93/13123, EP-A 0 558 416, EP-A 0 471 612, EP-A 0 384 842, EP-B 0 097 572, WO/99 25725.

In addition, the steroid derivatives that are known from EP 0 138 504 B1 can be mentioned. The 7α -[9-(4,4,5,5,5-pentafluoropentylsulfinyl)-n-nonyl]-estra-1,3,5(10)-triene-3,17 β -diol is currently under clinical development for hormone-dependent tumors (breast cancer).

Pharmaceutical compositions that contain sex steroid inhibitors and that have a steroidal skeleton that has a 7α -side chain in the case of the simultaneous presence of at least one other substituent in 14-, 15- or 16-position are the subject matter of EP-A 0 376 576.

Antiestrogenically active estratrienes that can carry an 11 β -fluorine atom and carry an α -position side chain in 7-position, which has an amino group and a sulfur group and that is functionalized in the terminal position, are described in WO 98/07740.

The compounds according to the invention are compounds with stronger antiestrogenic action after peroral administration.

The antiuterus growth test in infant rats, s.c. and p.o. (test on antiestrogenic action in-vivo) confirms the antiestrogenic action of the compounds according to the invention. The test is performed as described below:

Uterus Growth Test in Infant Rats (Antiestrogenic Action)

Principle of the Method

In rodents, the uterus reacts to the administration of estrogens with an increase in weight (both proliferation and water retention). This growth can be inhibited in a dose-dependent manner by simultaneous administration of compounds that have an antiestrogenic action.

Execution of the Test

Animals:

Infant female rats weighing 35-45 g at the beginning of the test, 5-6 animals per dose.

Formulation and Administration of the Substances:

For the p.o. administration, the substances are dissolved in 1 part ethanol (E) and made up with 9 parts peanut oil (EÖ).

Test Batch

The young rats just dropped by the mothers are delivered for acclimation one day before the beginning of treatment and immediately supplied with food -- right in the cage. The treatment is then carried out once daily over 3 days in combination with 0.5 μ g of estradiol benzoate (EB). EB is always administered subcutaneously (s.c.), while the test substance is administered p.o. (perorally). 24 hours after the last administration, the animals are weighed, killed and the uteri are

202059-62481001

removed. The moist weight (less contents) is determined from the prepared uteri.

Controls

Negative control: Vehicle (E/EÖ), 0.2 ml/animal/day

Positive control: 0.5 µg of EB/0.1 ml/animal/day

Evaluation

The average values with standard deviation ($X + SD$) and the significance of the differences in the control group (EB) in the Dunnett Test ($p < 0.05$) are determined for each group from the relative organ weights (mg/100 g of body weight). The calculation of the inhibition (in %) compared to the EB-control is carried out with a program. The relative actions of the test substances are determined by co-variance analysis and regression analysis.

As pure antiestrogens for the purposes of this invention, those compounds of general formula I that show no action or, in the best case, only slightly agonistic action, in the in-vitro test on estrogenic action can be considered.

By means of the method described below, the estrogenic effect of the compounds according to the invention on the bones can be determined. In the case of selectively estrogenically active compounds, protective effects on the bones are observed with comparable dosages, while on the uterus, no stimulation, or in the best case, only slight stimulation, is noted.

Bone Studies

Method

Female rats that are three months old are ovariectomized and treated once daily with the test compound immediately after the operation for 28 days. The administration is carried out subcutaneously in castor oil/benzyl benzoate or arachis oil/ethanol. The animals are sacrificed on the day after the last administration, and femurs, tibia as well as the uteri are removed. The uteri are weighed, mounted and worked up for histological studies. The determination of the bone density is carried out ex vivo on prepared long bones via pQCT (Quantitative Computer Tomography). The measurements are made at a distance of 5-7 mm from the ball of the joint at the distal femur or the proximal tibia.

As an alternative, the action on the bones by measuring out the trabecular bone surface area of the secondary spongiosa on histologic preparations of the distal femur or the proximal tibia is noted. The result is expressed as the proportion, in percent, of the trabecular bone surface area to the measured total bone surface area (TB/BV). The bone density that is measured via QCT and the trabecular bone surface area that is determined at the histologic section correlate well with one another. A comparison of the two measurement variables is therefore permissible.

The transition between the pure antiestrogens and the partial agonists, the tissue-selective estrogens, is seamless. Compounds that have a slightly agonistic action can also be used

202050-62481001

in the indications that are mentioned below for pure antiestrogens.

The compounds according to the invention, especially if they are pure antiestrogens, are suitable for treatment of estrogen-dependent diseases, for example breast cancer (second-line treatment of tamoxifen-resistant breast cancer; for adjuvant treatment of breast cancer instead of tamoxifen), endometrial cancer, prostate cancer, prostatic hyperplasia, anovulatory infertility and melanoma.

In addition, the pure antiestrogens of general formula I can be used as components in the products that are described in EP 346 014 B1, which contain an estrogen and a pure antiestrogen, specifically for simultaneous, sequential or separate use for selective estrogen therapy of peri- or postmenopausal women. The compounds of general formula I, especially if these are pure antiestrogens, can be used together with antigestagens (competitive progesterone antagonists) for the treatment of hormone-dependent tumors (EP 310 542 A).

Other indications in which the compounds of the general formula can be used are male hair loss, diffuse alopecia, alopecia that is caused by chemotherapy as well as hirsutism (Hye-Sun Oh and Robert C. Smart, Proc. Natl. Acad. Sci. USA, 93 (1996) 12525-12530).

In addition, the compounds of general formula I can be used for the production of medications for treating endometriosis and endometrial carcinomas.

The compounds of general formula I can also be used for the production of pharmaceutical compositions for male and female birth control (male birth control: DE-A 195 10 862.0).

The compounds of general formula I with tissue-selective partial estrogenic action can be used primarily for prophylaxis and treatment of osteoporosis and for the production of preparations for substitution therapy in pre-, peri- and post-menopause (HRT = hormone replacement therapy) (Black, L. J.; Sato, M.; Rowley, E. R.; Magee, D. E.; Bekele, A.; Williams, D. C.; Cullinan, G. J.; Bendele, R.; Kauffman, R. F.; Bensch, W. R.; Frolik, C. A.; Termine, J. D. and Bryant, H. U.: Raloxifene [LY 139481 HCl] Prevents Bone Loss and Reduces Serum Cholesterol without Causing Uterine Hypertrophy in Ovariectomized Rats; J. Clin. Invest. 93: 63-69, 1994).

The invention also relates to pharmaceutical preparations that contain at least one compound of general formula I (or physiologically compatible addition salts with organic and inorganic acids of them) and the use of these compounds for the production of pharmaceutical agents, especially for treating estrogen-dependent diseases and tumors and pharmaceutical agents for hormone replacement therapy (HRT).

The compounds according to the invention and the acid addition salts are suitable for the production of pharmaceutical compositions and preparations. As active ingredients, the pharmaceutical compositions or pharmaceutical agents contain one or more of the compounds according to the invention or their acid addition salts, optionally mixed with other pharmacologically or

202050 6484001

pharmaceutically active substances. The production of the pharmaceutical agents is carried out in a known way, whereby the known and commonly used pharmaceutical adjuvants and other commonly used vehicles and diluents can be used.

As such vehicles and adjuvants, for example, those are suitable that are recommended or indicated in the following bibliographic references as adjuvants for pharmaceuticals, cosmetics and related fields: Ullmans Encyklopädie der technischen Chemie [Ullman's Encyclopedia of Technical Chemistry], Volume 4 (1953), pages 1 to 39; Journal of Pharmaceutical Sciences, Volume 52 (1963), pages 918 and ff.; issued by Czetsch-Lindenwald, Hilfsstoffe für Pharmazie und angrenzende Gebiete [Adjuvants for Pharmaceuticals and Related Fields]; Pharm. Ind. Issue 2, 1961, pages 72 and ff.; Dr. H. P. Fiedler, Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete [Dictionary of Adjuvants for Pharmaceuticals, Cosmetics and Related Fields] Cantor KG, Aulendorf in Württemberg 1971.

The compounds can be administered orally or parenterally, for example intraperitoneally, intramuscularly, subcutaneously or percutaneously. The compounds can also be implanted in the tissue. The amount of the compounds to be administered varies within a wide range and can cover any effective amount. Based on the condition to be treated and the type of administration, the amount of the administered compound can be 0.1-25 mg/kg of body weight, preferably 0.5-5 mg/kg of body weight, per day. In humans, this corresponds to a daily dose of 5 to 1250 mg. The

preferred daily dosage in humans is 50 to 200 mg. This is true especially for tumor therapy.

For oral administration, capsules, pills, tablets, coated tablets, etc., are suitable. In addition to the active ingredient, the dosage units can contain a pharmaceutically compatible vehicle, such as, for example, starch, sugar, sorbitol, gelatin, lubricant, silicic acid, talc, etc. The individual dosage units for oral administration can contain, for example, 5 to 500 mg of active ingredient.

To achieve better bio-availability of the active ingredient, the compounds can also be formulated as cyclodextrin clathrates. For this purpose, the compounds are reacted with α -, β - or γ -cyclodextrin or derivatives thereof (PCT/EP95/02656).

For parenteral administration, the active ingredients can be dissolved or suspended in a physiologically compatible diluent. As diluent, very frequently oils with or without the addition of a solubilizer, a surfactant, a suspending agent or emulsifier are used. Examples of oils that are used are olive oil, peanut oil, cottonseed oil, soybean oil, castor oil and sesame oil.

The compounds of general formula I can also be formulated in the form of a solution that is determined for oral administration and that in addition to the active compound of general formula I contains

- a) a pharmaceutically compatible oil and/or
 - b) a pharmaceutically compatible lipophilic surfactant
- and/or

- c) a pharmaceutically compatible hydrophilic surfactant and/or
- d) a pharmaceutically compatible water-miscible solvent.

In this respect, reference is made in addition to WO 97/21440.

The compounds can also be used in the form of a depot injection or an implant preparation, which can be formulated in such a way that a delayed release of active ingredient is made possible.

As inert materials, implants can also contain, for example, biodegradable polymers or synthetic silicones such as, for example, silicone gum. In addition, the active ingredients can be embedded in, for example, a patch for percutaneous administration.

For the production of intravaginal systems (e.g., vaginal rings) or intrauterine systems (e.g., pessaries, spirals) that are loaded with active compounds of general formula I, various polymers such as, for example, silicone polymers, ethylene vinyl acetate, polyethylene or polypropylene are suitable.

The compounds according to the invention can be produced as described below. The examples below are used for a more detailed explanation of the invention. Other compounds of general formula I can be obtained by an analogous procedure using analogous reagents in the data contained in the examples.

Side chains R^{11} that do not contain any sulfur groups can be created analogously to the corresponding 7α -position side chains of the compounds that are described in PCT/EP98/08470, whereby

the 11 β -(5-chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol that is described here in Example 1d or the 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol that is described in Example 3a is now to be taken as a starting material.

A thio bridge in the side chain can be oxidized with sodium periodate to form sulfoxide; the sulfones are obtained from the sulfides with a peracid as an oxidizing agent, e.g., m-chloroperbenzoic acid.

The saponification of the ester groupings as well as esterification and etherification of free hydroxy groups is carried out in each case according to established processes of organic chemistry. By observing the varied reactivity of the esterified and free 3- and 17-hydroxy groups, the 3,17-diesters can be cleaved selectively in 3-position, and the 3-hydroxy-17-acyloxy compound can then be additionally functionalized specifically in the 3-position; it is equally possible to esterify or to etherify the 3,17-dihydroxy compound selectively only in the 3-position and then to introduce specifically another radical into the 17-position as already in the 3-position.

The acid addition salts of the compounds of general formula I can also be produced from the compounds of general formula I according to standard processes.

The examples below are used for a more detailed explanation of the invention:

10018429-050700

Example 1

11 β -[5-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

a) 11 β -[5-(*tert*-Butyldimethylsilyloxy)pentyl]-3,3-(2,2-dimethyltrimethylenedioxy)-5 α -estr-9-ene-5,17 β -diol

1.82 g of magnesium in 15 ml of absolute tetrahydrofuran is reacted under nitrogen with a solution of 21.1 g of 1-bromo-5-*tert*-butyldimethylsilyloxypentane [Tetrahedron Letters 1982, 4147-4150] in 40 ml of absolute tetrahydrofuran to form a Grignard reagent. At 0°C, 0.25 g of copper(I) chloride is added, and it is stirred for 15 more minutes, before a solution of 9.36 g of 3,3-(2,2-dimethyltrimethylenedioxy)-5,10 α -epoxy-5 α -estr-9(11)-en-17 β -ol [Neef, G. et al., Tetrahedron, (1993), 49, pp. 833-840]] in 40 ml of absolute tetrahydrofuran is added in drops, such that the internal temperature does not exceed 8°C. After the addition is completed, it is stirred for 90 more minutes while being cooled in an ice bath. For working-up, the reaction mixture is added to saturated ammonium chloride solution while being stirred, extracted three times with ethyl acetate, the combined ethyl acetate phases are washed with saturated sodium chloride solution, dried on sodium sulfate and evaporated to the dry state in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant yields 5.02 g of 11 β -[5-(*tert*-butyldimethylsilyloxy)pentyl]-3,3-(2,2-dimethyltrimethylenedioxy)-5 α -estr-9-ene-5,17 β -diol as a foam.

b) 17 β -Hydroxy-11 β -(5-hydroxypentyl)estra-4,9-dien-3-one

4.96 g of 11 β -[5-(*tert*-butyldimethylsilyloxy)pentyl]-3,3-(2,2-dimethyltrimethylenedioxy)-5 α -estr-9-ene-5,17 β -diol in 36 ml of tetrahydrofuran is stirred under nitrogen with 40 ml of glacial acetic acid and 20 ml of water for 3 hours at a bath temperature of 50°C. Then, it is carefully poured onto saturated sodium bicarbonate solution, extracted three times with ethyl acetate, the combined organic phases are washed with saturated sodium chloride solution, dried on sodium sulfate and evaporated to the dry state in a vacuum. By preparative column chromatography on silica gel with hexane/acetone as an eluant, 3.57 g of 17 β -hydroxy-11 β -(5-hydroxyphenyl)estra-4,9-dien-3-one is obtained as a foam.

c) 11 β -(5-Hydroxypentyl)-9 ξ -estra-1,3,5(10)-triene-3,17 β -diol

A solution of 1.08 g of 17 β -hydroxy-11 β -(5-hydroxypentyl)estra-4,9-dien-3-one in 19 ml of ethanol is refluxed with 0.19 g of palladium on activated carbon for 20 hours at a bath temperature of 100°C. After cooling, it is filtered on Celite, rewashed with ethyl acetate, the filtrate is evaporated to the dry state and chromatographed on silica gel with hexane/acetone. Recrystallization from methylene chloride results in 0.46 g of 11 β -(5-hydroxypentyl)-9 ξ -estra-1,3,5(10)-triene-3,17 β -diol with a melting point of 152°C.

202050-6481001

d) 11 β -(5-Chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol

426 mg of 11 β -(5-hydroxypentyl)-9 ξ -estra-1,3,5(10)-triene-3,17 β -diol is suspended in 7.2 ml of carbon tetrachloride and 2.4 ml of acetonitrile, mixed with 682 mg of triphenyl-phosphine and stirred for 90 minutes at room temperature. For working-up, the reaction solution is mixed with methylene chloride, washed with saturated sodium bicarbonate solution and saturated sodium chloride solution, dried on sodium sulfate, and concentrated by evaporation in a vacuum. By preparative column chromatography on silica gel with methylene chloride/ethyl acetate, the 9 β -isomers can be separated. Subsequent crystallization from methylene chloride yields 238 mg of 11 β -(5-chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol with a melting point of 159°C.

e) 11 β -[5-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

189 mg of 11 β -(5-chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 3 ml of dried dimethylformamide, mixed with 598 mg of methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}amine [DE 196 35 525.7] and stirred for 24 hours at 100°C under nitrogen. After the reaction mixture is cooled, it is diluted with ethyl acetate, washed once with saturated sodium bicarbonate solution and twice with saturated sodium chloride solution, dried on sodium sulfate and concentrated by evaporation in a vacuum. After preparative column chromatography on silica gel with ethyl acetate/methanol as an eluant, 222 mg of 11 β -[5-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}-

amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam $[\alpha]_D^{22} = +72.1^\circ$ ($c = 0.172$ in chloroform).

Example 2

11 β -(5-{3-[(4,4,5,5,5-Pentafluoropentyl)sulfanyl]-propylamino}pentyl)estra-1,3,5(10)-triene-3,17 β -diol

98 mg of 11 β -(5-chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 1.5 ml of dried dimethylformamide, mixed with 3 ml of 3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propylamine and stirred for 16 hours under nitrogen at a bath temperature of 80°C. After the reaction solution is cooled, it is diluted with ethyl acetate, washed once with water and twice with saturated sodium chloride solution, dried on sodium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol with the addition of ammonia results in 31 mg of 11 β -(5-{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propylamino}pentyl)-estra-1,3,5(10)-triene-3,17 β -diol as a foam.

Example 3

11 β -[5-(Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl)-amino}pentyl]estra-1,3,5(10)-triene-3,17 β -diol

a) 11 β -(5-Iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol

1.46 g of 11 β -(5-chloropentyl)estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 24 ml of ethylmethylketone, mixed with 1.82 g of sodium iodide and stirred overnight at a bath temperature of 90°C. For working-up, the reaction mixture is

added to water, extracted three times with ethyl acetate, washed with sodium thiosulfate solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 1.86 g of 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol is obtained as a crude product, which is used in the next reaction without further purification.

b) 11 β -[5-(Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amino)-pentyl]estra-1,3,5(10)-triene-3,17 β -diol

1.0 g of 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol and 2.47 g of methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amine are dissolved in 21 ml of N-methylpyrrolidone and stirred for 3 hours at 80°C. After the reaction solution is cooled to room temperature, the batch is added to semi-saturated sodium chloride solution, extracted three times with diethyl ether, the combined organic phases are washed with water and saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 0.69 g of 11 β -[5-(methyl{3-[(2-pyridyl-methyl)sulfanyl]propyl}-amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = +62.2^\circ$ (c = 0.519 in chloroform).

10018499-050709
202050-64481001

Example 4

11 β -[5-(Methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}amino)-
pentyl]estra-1,3,5(10)-triene-3,17 β -diol

250 mg of 11 β -[5-(methyl{3-[(2-pyridylmethyl)sulfanyl]-
propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol is
dissolved in 8.1 ml of methanol and 0.44 ml of water, mixed with
161 mg of sodium metaperiodate and stirred for 16 hours at room
temperature. For working-up, the reaction mixture is added to
semi-saturated sodium chloride solution, extracted three times
with methylene chloride, the combined organic phases are dried on
magnesium sulfate and concentrated by evaporation in a vacuum.
Preparative column chromatography on silica gel with methylene
chloride/methanol as an eluant yields 53 mg of 11 β -[5-(methyl{3-
[(2-pyridylmethyl)sulfinyl]propyl}-amino)pentyl]estra-1,3,5(10)-
triene-3,17 β -diol as a foam.

$[\alpha]_D^{22} = +58.9^\circ$ (c = 0.501 in chloroform).

Example 5

11 β -[5-(Methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-
propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

839 mg of 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -
diol and 1.42 g of methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-
propyl}amine are dissolved in 18 ml of N-methylpyrrolidone and
stirred for 90 minutes at a bath temperature of 80°C. After the
reaction solution is cooled to room temperature, the batch is
added to semi-saturated sodium chloride solution, extracted three
times with diethyl ether, the combined organic phases are washed

202050 6481001

with water and saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 0.94 g of 11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = +75.0^\circ$ (c = 0.5 in chloroform).

Example 6

11 β -[5-(Methyl{3-[4-(trifluoromethyl)benzylsulfinyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

630 mg of 11 β -[5-(methyl{3-[4-(trifluoromethyl)benzyl-sulfanyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 17.4 ml of methanol and 0.95 ml of water, mixed with 329 mg of sodium metaperiodate and stirred at room temperature for 16 hours. For working-up, the reaction mixture is added to semi-saturated sodium chloride solution, extracted three times with methylene chloride, the combined organic phases are dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 216 mg of 11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = +54.1^\circ$ (c = 0.501 in chloroform).

10018429-050702

Example 7

11 β -{5-[(2S)-2-{[4-(Trifluoromethyl)phenyl]sulfanylmethyl}-pyrrolidin-1-yl]pentyl}estra-1,3,5(10)-triene-3,17 β -diol

820 mg of 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol and 458 mg of (S)-2-{[4-(trifluoromethyl)phenyl]sulfanylmethyl}pyrrolidine are dissolved in 16 ml of N-methylpyrrolidone and stirred for 3 hours at a bath temperature of 90°C. After the reaction solution is cooled to room temperature, the batch is added to semi-saturated sodium chloride solution, extracted three times with ethyl acetate, the combined organic phases are washed three times with water, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 561 mg of 11 β -{5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfanylmethyl}pyrrolidin-1-yl]pentyl}estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = +33.1^\circ$ (c = 0.5195 in chloroform).

Example 8

11 β -{5-[(2S)-2-{[4-(Trifluoromethyl)phenyl]sulfinylmethyl}-pyrrolidin-1-yl]pentyl}estra-1,3,5(10)-triene-3,17 β -diol

975 mg of 11 β -(5-iodopentyl)estra-1,3,5(10)-triene-3,17 β -diol and 540 mg of (S)-2-{[4-(trifluoromethyl)phenyl]sulfinylmethyl}pyrrolidine are dissolved in 19 ml of N-methylpyrrolidone and stirred for 3 hours at a bath temperature of 90°C. After the reaction solution is cooled to room temperature, the batch is added to semi-saturated sodium chloride solution, extracted three

times with ethyl acetate, the combined organic phases are washed twice with water, dried on magnesium sulfate, and concentrated by evaporation in a vacuum. Chromatography on silica gel with methylene chloride/methanol yields 222 mg of 11 β -{5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfinyl-methyl}pyrrolidine-1-yl]pentyl}-estra-1,3,5(10)-triene-3,17 β -diol as a foam. $[\alpha]_D^{22} = +42.6^\circ$ (c = 0.5145 in chloroform).

Example 9

11 β -{4-[2-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

a) 11 β -{4-[2-(*tert*-Butyldimethylsilyloxy)ethyl]phenyl}-3,3-(2,2-dimethyltrimethylene-dioxy)-5 α -estr-9-ene-5,17 β -diol

1.94 g of magnesium chips is introduced under nitrogen into 15 ml of absolute tetrahydrofuran and mixed with about 2.5 g of 1-bromo-4-[2-(*tert*-butyldimethylsilyloxy)ethyl]benzene in 4 ml of absolute tetrahydrofuran as well as a spatula tip full of iodine. After the reaction is started, the residual 22.8 g of 1-bromo-4-[2-(*tert*-butyl-dimethylsilyloxy)ethyl]benzene is added in drops to 36 ml of absolute tetrahydrofuran, and after the addition is completed, it is stirred for 1 hour at 80°C. Then, it is cooled to 0°C, mixed with 267 mg of copper(I) chloride and stirred for 30 more minutes under cold conditions. Then, 10 g of 3,3-(2,2-dimethyltrimethylenedioxy)-5,10 α -epoxy-5 α -estr-9(11)-en-17 β -ol is added in drops while being cooled in an ice bath in such a way that the internal temperature does not exceed +8°C. After 2 hours of stirring at 0°C, the reaction mixture is added to

204050-6248T001

saturated ammonium chloride solution, extracted three times with ethyl acetate, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant results in 11.97 g of 11 β -{4-[2-(*tert*-butyldimethylsilyloxy)ethyl]phenyl}-3,3-(2,2-dimethyltrimethylenedioxy)-5 α -estr-9-ene-5,17 β -diol.

b) 17 β -Hydroxy-11 β -[4-(2-hydroxyethyl)phenyl]estra-4,9-dien-3-one

11.27 g of 11 β -{4-[2-(*tert*-butyldimethylsilyloxy)ethyl]phenyl}-3,3-(2,2-dimethyl-trimethylenedioxy)-5 α -estr-9-ene-5,17 β -diol is dissolved in 70 ml of tetrahydrofuran, mixed with 86 ml of glacial acetic acid as well as 43 ml of water and stirred for 1 hour at a bath temperature of 50°C. After the cooling, the reaction solution is carefully added to ice-cold sodium bicarbonate solution, extracted three times with ethyl acetate, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. After preparative column chromatography on silica gel with hexane/acetone as an eluant, 6.73 g of 17 β -hydroxy-11 β -[4-(2-hydroxyethyl)phenyl]estra-4,9-dien-3-one is obtained as a foam.

c) 11 β -[4-(2-Chloroethyl)phenyl]-17 β -hydroxyestra-4,9-dien-3-one

6.32 g of 17 β -hydroxy-11 β -[4-(2-hydroxyethyl)phenyl]estra-4,9-dien-3-one is dissolved in 160 ml of methylene chloride, mixed with 7.65 g of triphenylphosphine and 16.1 ml of perchloro-

202050-62481001

acetone and stirred for 1 hour at room temperature. Then, the reaction mixture is concentrated by evaporation in a vacuum and chromatographed on silica gel with hexane/ethyl acetate as well as methylene chloride/methanol as an eluant. 3.19 g of 11 β -[4-(2-chloroethyl)phenyl]-17 β -hydroxyestra-4,9-dien-3-one is obtained as a foam.

d) 11 β -[4-(2-Chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol

A solution of 2.94 g of 11 β -[4-(2-chloroethyl)phenyl]-17 β -hydroxyestra-4,9-dien-3-one in 70 ml of ethanol is refluxed with 458 mg of palladium on activated carbon for 5 hours at a bath temperature of 100°C. After cooling to room temperature, it is filtered on Celite, rewashed with ethanol and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant yields 522 mg of 11 β -[4-(2-chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol as a foam.

e) 11 β -[4-(2-Iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol

312 mg of 11 β -[4-(2-chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 5 ml of ethylmethylketone, mixed with 341 mg of sodium iodide and stirred for 16 hours at 90°C. For working-up, the reaction mixture is added to water, extracted three times with ethyl acetate, washed with thiosulfate solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 382 mg of 11 β -[4-(2-iodoethyl)phenyl]-

estra-1,3,5(10)-triene-3,17 β -diol is obtained as a crude product, which is used in the next stage without further purification.

f) 11 β -{4-[2-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

382 mg of 11 β -[4-(2-iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol and 603 mg of methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propyl}amine are dissolved in 7.5 ml of N-methylpyrrolidone and stirred for 150 minutes at a bath temperature of 90°C. After the reaction solution is cooled to room temperature, the batch is added to dilute sodium chloride solution, extracted three times with diethyl ether, the combined organic phases are washed with water and saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 180 mg of 11 β -{4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = -14.4^\circ$ (c = 0.502 in chloroform).

Example 10

11 β -{4-[2-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfinyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

205 mg of 11 β -[4-(2-chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol and 633 mg of methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfinyl]propyl}amine are dissolved in 5 ml of dimethylformamide, and stirred for 5 hours at a bath temperature

20250162481001

of 90°C and for 16 hours at room temperature. For working-up, the batch is added to semi-saturated sodium chloride solution, extracted three times with ethyl acetate, washed twice with water, dried on magnesium sulfate and concentrated by evaporation in a vacuum. The residue is chromatographed on silica gel with methylene chloride/methanol as an eluant. 54 mg of 11 β -{4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfinyl]propyl}-(amino)ethyl]-phenyl}estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam, $[\alpha]_D^{22} = -12.9^\circ$ (c = 0.509 in chloroform).

Example 11

11 β -{4-[2-(Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amino)-ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

a) 11 β -[4-(2-Hydroxyethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol

A solution of 11.7 g of 17 β -hydroxy-11 β -[4-(2-hydroxyethyl)phenyl]estra-4,9-dien-3-one in 295 ml of methanol is stirred with 8.54 g of palladium hydroxide on activated carbon (20%) and 2.16 g of magnesium oxide for 5 hours at a bath temperature of 70°C. After cooling, it is filtered on Celite, washed with methanol and concentrated by evaporation in a vacuum. The residue is chromatographed on silica gel with hexane/acetone as an eluant. 4.54 g of 11 β -[4-(2-hydroxyethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam, $[\alpha]_D^{22} = -175.5^\circ$ (c = 0.502 in pyridine).

10018439-060792

b) 11 β -[4-(2-Chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol

4.50 g of 11 β -[4-(2-hydroxyethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 114 ml of methylene chloride and mixed in succession with 5.45 g of triphenylphosphine and 11.5 ml of perchloroacetone. Then, it is stirred for 90 minutes at room temperature. For working-up, the reaction mixture is concentrated by evaporation in a vacuum and chromatographed on silica gel with methylene chloride/methanol as an eluant. 4.02 g of 11 β -[4-(2-chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam, $[\alpha]_D^{22} = +10.7^\circ$ (c = 0.503 in chloroform).

c) 11 β -[4-(2-Iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol

411 mg of 11 β -[4-(2-chloroethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 6.5 ml of ethylmethylketone, mixed with 449 mg of sodium iodide and stirred for 16 hours at 90°C. For working-up, the reaction mixture is added to dilute sodium chloride solution, extracted three times with ethyl acetate, washed with thiosulfate solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 479 mg of 11 β -[4-(2-iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol is obtained as a crude product, which is used in the next stage without further purification.

d) 11 β -{4-[2-(Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amino)-ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

470 mg of 11 β -[4-(2-iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol and 554 mg of methyl{3-[(2-pyridylmethyl)sulfanyl]-propyl}amine are dissolved in 9 ml of N-methylpyrrolidone and stirred for 2 hours at 90°C. After the reaction solution is cooled to room temperature, the batch is added to semi-saturated sodium chloride solution, extracted three times with diethyl ether, the combined organic phases are washed with water, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 93 mg of 11 β -{4-[2-(methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amino)-ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = -31.8^\circ$ (c = 0.513 in chloroform).

Example 12

11 β -{4-[2-(Methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}amino)-ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

68 mg of 11 β -{4-[2-(methyl{3-[(2-pyridylmethyl)sulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 2 ml of methanol and 0.12 ml of water, mixed with 28 mg of sodium metaperiodate and stirred for 16 hours at room temperature. For working-up, the reaction mixture is added to semi-saturated sodium chloride solution, extracted three times with methylene chloride, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by

1004949-6443001

evaporation in a vacuum. The residue is purified by means of preparative thin-layer chromatography with methylene chloride/methanol with the addition of ammonia as a mobile solvent. 31 mg of 11 β -{4-[2-(methyl{3-[(2-pyridylmethyl)-sulfinyl]propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam, $[\alpha]_D^{22} = -33.3^\circ$ (c = 0.51 in chloroform).

Example 13

11 β -{4-[2-(Methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

466 mg of 11 β -[4-(2-iodoethyl)phenyl]estra-1,3,5(10)-triene-3,17 β -diol and 359 mg of methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amine are dissolved in 9 ml of N-methylpyrrolidone and stirred for 4 hours at a bath temperature of 40°C. After the reaction solution is cooled to room temperature, the batch is added to semi-saturated sodium chloride solution, extracted three times with diethyl ether, the combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with methylene chloride/methanol as an eluant yields 256 mg of 11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}-amino)ethyl]phenyl}-estra-1,3,5(10)-triene-3,17 β -diol as a foam, $[\alpha]_D^{22} = -20.2^\circ$ (c = 0.505 in chloroform).

10018439-050702

Example 14

11 β -{4-[2-(Methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}-amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

100 mg of 11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}amino)ethyl]-phenyl}estra-1,3,5(10)-triene-3,17 β -diol is dissolved in 2.6 ml of methanol and 0.15 ml of water, mixed with 36 mg of sodium metaperiodate, and stirred for 4 hours at room temperature. For working-up, the reaction mixture is added to semi-saturated sodium chloride solution, extracted three times with methylene chloride, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. The residue is purified by means of preparative thin-layer chromatography with methylene chloride/methanol with the addition of ammonia as a mobile solvent. 48 mg of 11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}amino)ethyl]-phenyl}estra-1,3,5(10)-triene-3,17 β -diol is obtained as a foam, $[\alpha]_D^{22} = -28.8^\circ$ (c = 0.511 in chloroform).

Production of the Reagents

3-[(4,4,5,5,5-Pentafluoropentyl)sulfanyl]propylamine

25 g of S-(4,4,5,5,5-pentafluoropentyl)thioacetate [Singh, S. M. et al., Tet. Lett., (1994), 35, pp. 9141-9144] is introduced at 0°C into 250 ml of absolute acetonitrile and mixed drop by drop with 39.2 ml of a 30% sodium methylate solution. Stirring is continued for 15 minutes under cold conditions, before 23.2 g of 3-bromopropylamine hydrobromide is introduced in

202050: 62494001

portions. Then, it stirred for one more hour at 0°C. The reaction mixture is then added to water, extracted three times with diethyl ether, washed neutral, dried on magnesium sulfate and concentrated by evaporation. After preparative column chromatography on silica gel with methylene chloride/methanol with the addition of ammonia as an eluant, 22.54 g of 3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]propylamine is obtained.

Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amine

a) S-(2-pyridylmethyl)thioacetate

10.0 g of 2-(Chloromethyl)pyridine hydrochloride is introduced into 100 ml of acetone, mixed with 14.0 g of potassium thioacetate and refluxed for 2 hours at 80°C. For working-up, the reaction mixture is diluted with ethyl acetate and added to water. It is extracted three times with ethyl acetate, the combined organic phases are washed with saturated sodium bicarbonate solution and with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. After preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant, 9.78 g of S-(2-pyridylmethyl)thioacetate is obtained.

b) 2-[(3-Chloropropyl)sulfanylmethyl]pyridine

9.78 g of S-(2-pyridylmethyl)thioacetate in 90 ml of absolute methanol is mixed drop by drop with 10.9 ml of a 30% methanolic sodium methylate solution while being cooled in an ice bath, and after the addition is completed, it is stirred for 20

more minutes. Then, 8.6 ml of 1-bromo-3-chloropropane is added in drops at 4°C and stirred for two more hours under cold conditions and for one more hour at room temperature. Then, the reaction mixture is added to water, extracted three times with ethyl acetate, the combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant yields 11.7 g of 2-[(3-chloropropyl)sulfanylmethyl]-pyridine.

c) Methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}amine

5.85 g of 2-[(3-chloropropyl)sulfanylmethyl]pyridine in 33.5 ml of tetrahydrofuran is mixed with 11.7 g of sodium iodide in a pressure vessel. Then, 17.05 g of methylamine is condensed at -20°C and heated overnight in a pressure vessel at a bath temperature of 50°C. After the reaction vessel was opened at -20°C, it is allowed to come to room temperature to allow excess methylamine to evaporate. The reaction solution is added to dilute sodium chloride solution, extracted three times with methylene chloride, the combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 5.7 g of methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}-amine is obtained as a crude product.

Methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amine

a) 1-[(3-Bromopropyl)sulfanylmethyl]-4-(trifluoromethyl)benzene

25.0 g of 4-(trifluoromethyl)benzyl bromide is introduced into 84 ml of acetonitrile, mixed with 8.3 ml of trimethylene sulfide and stirred for 48 hours at room temperature. Then, the reaction mixture is evaporated to the dry state and chromatographed on silica gel with hexane/methylene chloride. 29.47 g of 1-[(3-bromopropyl)sulfanyl-methyl]-4-(trifluoromethyl)benzene is obtained as an oil.

b) Methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amine

18.8 g of methylamine is condensed at -20°C in a solution of 10.0 g of 1-[(3-bromopropyl)sulfanylmethyl]-4-(trifluoromethyl)-benzene in 37.0 ml of tetrahydrofuran, and it is stirred overnight at room temperature in a pressure vessel. After the pressure vessel was opened at -20°C, it is allowed to come to room temperature to allow excess methylamine to evaporate. The reaction solution is added to dilute sodium chloride solution, extracted three times with methylene chloride, the combined organic phases are washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 8.32 g of methyl{3-[4-(trifluoromethyl)benzylsulfanyl]propyl}amine is obtained as a crude product.

202050-62431001

1-Bromo-4-[2-(*tert*-butyldimethylsilyloxy)ethyl]benzene

70 g of 2-(4-bromophenyl)ethanol is dissolved in 350 ml of absolute tetrahydrofuran and mixed with 4.97 g of imidazole. Then, 55 g of *tert*-butyldimethylsilyl chloride in 230 ml of absolute tetrahydrofuran is added in drops and stirred for 90 minutes at room temperature. For working-up, the reaction mixture is added to ice water, extracted three times with diethyl ether, washed with water, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant yields 111 g of 1-bromo-4-[2-(*tert*-butyldimethylsilyloxy)ethyl]benzene as a clear oil.

(S)-2-{[4-(Trifluoromethyl)phenyl]sulfanylmethyl}pyrrolidine

a) (S)-*tert*-Butyl-2-{[4-(trifluoromethyl)phenyl]sulfanylmethyl}-pyrrolidine-1-carboxylate

1.93 ml of a 30% sodium methylate solution is slowly added in drops at 0°C under inert gas to a solution that consists of 1.4 g of 4-(trifluoromethyl)thiophenol in 15 ml of absolute acetone. After the addition is completed, it is stirred for 30 more minutes under cold conditions, before 1.85 g of (S)-*tert*-butyl-2-(bromomethyl)pyrrolidine-1-carboxylate [Katzenellenbogen, J. A. et al.; J. Med. Chem., (1994), 37, pp. 928-937] is added in drops to 5 ml of absolute acetone. Then, it is stirred for 30 more minutes under cold conditions and for 12 hours at room temperature. For working-up, the reaction mixture is added to water, extracted three times with diethyl ether, washed twice

with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. Preparative column chromatography on silica gel with hexane/ethyl acetate as an eluant yields 1.8 g of (S)-*tert*-butyl-2-{[4-(trifluoromethyl)-phenyl]sulfanylmethyl}pyrrolidine-1-carboxylate as an oil.

b) (S)-2-{[4-(Trifluoromethyl)phenyl]sulfanylmethyl}pyrrolidine
 920 mg of (S)-*tert*-butyl-2-{[4-(trifluoromethyl)phenyl]-sulfanylmethyl}pyrrolidine-1-carboxylate is added to a mixture, cooled to 0°C, that consists of 7.2 ml of trifluoroacetic acid, 0.22 ml of triisopropylsilane and 0.30 ml of water, and it is stirred for 1 hour at 0°C. Then, the reaction is set at pH 10 with 10% potassium hydroxide solution, extracted three times with methylene chloride, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 458 mg of (S)-2-{[4-(trifluoromethyl)-phenyl]sulfanylmethyl}-pyrrolidine is obtained as a crude product.

(S)-2-{[4-(Trifluoromethyl)phenyl]sulfinylmethyl}pyrrolidine

a) (S)-*tert*-Butyl-2-{[4-(trifluoromethyl)phenyl]sulfinylmethyl}-pyrrolidine-1-carboxylate

1.0 g of (S)-*tert*-butyl-2-{[4-(trifluoromethyl)phenyl]-sulfanylmethyl}pyrrolidine-1-carboxylate is dissolved in 48 ml of methanol and 2.6 ml of water, mixed with 833 mg of sodium metaperiodate and stirred for two days at room temperature and for one hour at 50°C. For working-up, the reaction mixture is

added to semi-saturated sodium chloride solution, extracted three times with methylene chloride, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. The residue is purified by preparative column chromatography with hexane/acetone as an eluant. 929 mg of (S)-*tert*-butyl-2-{[4-(trifluoromethyl)phenyl]-sulfinylmethyl}pyrrolidine-1-carboxylate is obtained.


b) (S)-2-{[4-(Trifluoromethyl)phenyl]sulfinylmethyl}pyrrolidine

920 mg of (S)-*tert*-butyl-2-{[4-(trifluoromethyl)phenyl]-sulfinyl-methyl}pyrrolidine-1-carboxylate is added to a mixture, cooled to 0°C, that consists of 7.2 ml of trifluoroacetic acid, 0.22 ml of triisopropylsilane and 0.30 ml of water, heated for a short time to room temperature until the educt is dissolved and then stirred for 30 minutes at 0°C. Then, the reaction is set at pH 10 with 10% potassium hydroxide solution, extracted three times with methylene chloride, washed with saturated sodium chloride solution, dried on magnesium sulfate and concentrated by evaporation in a vacuum. 549 mg of (S)-2-{[4-(trifluoromethyl)-phenyl]sulfinyl}pyrrolidine is obtained as a crude product.

202050-6448T001

Claims:

1. 11 β -Long-chain-substituted estratrienes of general formula I



in which

R^3 means a hydrogen atom, a hydrocarbon radical with up to 8 carbon atoms or a radical of partial formula $R^{3'}-C(O)-$, in which $R^{3'}$ means a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms or a phenyl radical,

R¹¹ means a radical of formula -A-B-Z-R²⁰,
in which

A stands for a direct bond, and

B stands for a straight-chain or branched-chain alkylene, alkenylene or alkinylene group with 4, 5 or 6 carbon atoms, or

A stands for a phenylene radical, and

B stands for a methylene, ethylene, propylene or trimethylene group, or

- A stands for a phenylenoxy radical, whereby the latter is bonded via a carbon atom to the 11-carbon atom of the steroid, and
- B stands for an ethylene group,
- Z stands for $-NR^{21}-$ and R^{21} stands for a C_1-C_3 alkyl group,

whereby R^{20} means

a hydrogen atom,
 a straight-chain or branched-chain alkyl,
 alkenyl or alkynyl group with up to 10 carbon atoms,
 whereby if A is a direct bond, R^{20} and R^{21} do not both simultaneously mean methyl, however, and, if A is a phenylenoxy radical, R^{20} and R^{21} do not both simultaneously mean methyl or ethyl, and if A is a phenylenoxy radical and B means an ethylene group, OR^{17b} should not be a hydroxy group and R^{17a} should not be a C_{1-4} alkyl group, and R^3 should not be a hydrogen atom,

or one of groupings

$-D-C_nF_{2n+1}$, whereby D is a straight-chain or branched-chain alkylene, alkenylene or alkynylene group with up to 8 carbon atoms and n is an integer from 1 to 8,
 D-aryl, whereby D has the already indicated meaning, and aryl stands for a phenyl, 1- or

10034439-050703

2-naphthyl radical or a heteroaryl radical that is optionally substituted in one or two places,

$-L-CH=CF-C_pF_{2p+1}$, whereby L is a straight-chain or branched-chain alkylene, alkenylene or alkynylene group with up to 7 carbon atoms and p is an integer from 1 to 7,

whereby in the three cases above in D or L, a methylene group can be replaced by a sulfur atom, a sulfone group or a sulfoxide group,

$-D-O-(CH_2)_q-aryl$, whereby D and aryl have the already indicated meanings, and q is 0, 1, 2 or 3,

$-D-O-(CH_2)_r-C_nF_{2n+1}$, whereby D and n have the already indicated meanings, and r stands for an integer from 1 to 5,

whereby in addition in all relevant cases above, R^{21} together with D with the inclusion of the nitrogen atom can then form a pyrrolidine ring that is substituted in 2- or 3-position,

or

if A is a direct bond or a phenylene radical, R^{20} and R^{21} with the nitrogen atom to which they are bonded form a saturated or unsaturated heterocyclic compound with 5 or 6

chain links, which optionally contains one or two additional heteroatoms, selected from nitrogen, oxygen and sulfur, and optionally is substituted,

whereby if A is a phenylene radical and B is a trimethylene radical, R^{21} and R^{20} do not form a methyl or ethyl group, or, together with the nitrogen atom to which they are bonded, do not form a pyrrolidine or piperidine ring,

and

R^{17a} in α - or β -position means a hydrogen atom, a C_{1-5} alkyl, a C_{2-5} alkenyl or a C_{2-5} alkynyl group or a trifluoromethyl group, or together with the radical OR 17b means a keto-oxygen atom, and

R^{17b} means a hydrogen atom or a radical of partial formula $R^{17'}--C(O)--$, in which $R^{17'}$ means a hydrogen atom or a hydrocarbon radical with up to 8 carbon atoms.

2. 11 β -Substituted estratrienes according to claim 1, in which R^3 is a hydrogen atom.

3. 11 β -Substituted estratrienes according to claim 1, in which R^3 is a benzoyl radical.

4. 11 β -Substituted estratrienes according to claim 1, in which R^{17b} is a hydrogen atom.

5. 11 β -Substituted estratrienes according to claim 1, in which R^{11} is selected from the group of the following side chains

202501543-0507001

$-(CH_2)_5N(CH_3)-(CH_2)_3-S-(CH_2)_3C_2F_5$
 $-(CH_2)_5NH-(CH_2)_3-S-(CH_2)_3C_2F_5$
 $-(CH_2)_5N(CH_3)-(CH_2)_3-S-CH_2-2-Pyridyl$
 $-(CH_2)_5N(CH_3)-(CH_2)_3-SO-CH_2-2-Pyridyl$
 $-(CH_2)_5N(CH_3)-(CH_2)_3-S-CH_2-p-CF_3-Phenyl$
 $-(CH_2)_5N(CH_3)-(CH_2)_3-SO-CH_2-p-CF_3-Phenyl$
 $-(CH_2)_5-[2-Pyrrolidin-1-yl]-CH_2-S-p-CF_3-Phenyl$
 $-(CH_2)_5-[2-Pyrrolidin-1-yl]-CH_2-SO-p-CF_3-Phenyl$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-S-(CH_2)_3C_2F_5$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-SO-(CH_2)_3C_2F_5$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-S-CH_2-2-Pyridyl$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-SO-CH_2-2-Pyridyl$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-S-CH_2-p-CF_3-Phenyl$
 $p\text{-Phenylene}-(CH_2)_2-N(CH_3)-(CH_2)_3-SO-CH_2-p-CF_3-Phenyl$

$-(CH_2)_5N(CH_3)(CH_2)_3C_2F_5$
 $-(CH_2)_5N(CH_3)(CH_2)_6C_2F_5$
 $-(CH_2)_5N(CH_3)(CH_2)_7C_2F_5$
 $-(CH_2)_5N(CH_3)(CH_2)_8C_2F_5$
 $-(CH_2)_6N(CH_3)(CH_2)_6C_2F_5$
 $-(CH_2)_6N(CH_3)(CH_2)_7C_2F_5$
 $-(CH_2)_6N(CH_3)(CH_2)_8C_2F_5$
 $-(CH_2)_5N(CH_3)(CH_2)_2C_4F_9$
 $-(CH_2)_5N(CH_3)(CH_2)_3C_6F_{13}$
 $-(CH_2)_5N(CH_3)(CH_2)_3C_8F_{17}$
 $-(CH_2)_5N(CH_3)(CH_2)_6C_4F_9$

$-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_6\text{C}_6\text{F}_{13}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_6\text{C}_8\text{F}_{17}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)\text{H}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_9\text{H}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CF}-\text{C}_2\text{F}_5$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CF}-\text{C}_3\text{F}_7$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CF}-\text{C}_5\text{F}_{11}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CF}-\text{C}_7\text{F}_{15}$
 $-(\text{CH}_2)_5-1\text{-Pyrrolidinyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{O}^i\text{Phenyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{O}^i\text{Benzyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{O}(\text{CH}_2)_5\text{C}_2\text{F}_5$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{CH}(\text{CH}_3)_2$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{-Pyridyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{-Phenyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_2\text{-p-Tolyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_2\text{-p-Ethoxyphenyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{-p-Tolyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{-p-Chlorophenyl}$
 $-(\text{CH}_2)_5\text{N}(\text{CH}_3)(\text{CH}_2)_3\text{-O-CH}_2\text{-Phenyl}$

[Key:]

$-(\text{CH}_2)_5\text{-[2-Pyrrolidin-1-yl]}-\dots = (\text{CH}_2)_5\text{-[2-pyrrolidine-1-yl]}-\dots$

$\text{p-Phenylene}-\dots = \text{p-phenylene}-\dots$

6. 11 β -Long-chain-substituted estratrienes of general formula I, namely

11 β -[5-(Methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propylamino}pentyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}-amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}-amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-propyl}amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}-amino)pentyl]estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfanyl-methyl}pyrrolidine-1-yl]pentyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-[(2S)-2-{[4-(trifluoromethyl)phenyl]sulfinyl-methyl}pyrrolidine-1-yl]pentyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -(4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfanyl]-propyl}amino)ethyl]phenyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -(4-[2-(methyl{3-[(4,4,5,5,5-pentafluoropentyl)sulfinyl]-propyl}amino)ethyl]phenyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -(4-[2-(methyl{3-[(2-pyridylmethyl)sulfanyl]propyl}-amino)ethyl]phenyl)estra-1,3,5(10)-triene-3,17 β -diol

11 β -(4-[2-(methyl{3-[(2-pyridylmethyl)sulfinyl]propyl}-amino)ethyl]phenyl)estra-1,3,5(10)-triene-3,17 β -diol

202050: 62481001

11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfanyl]-propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{4-[2-(methyl{3-[4-(trifluoromethyl)benzylsulfinyl]propyl}amino)ethyl]phenyl}estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(8,8,9,9,9-pentafluoro-nonyl)amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-nonyl-amino]pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(9,9,10,10,10-pentafluoro-decyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(8,8,9,9,9-pentafluoro-nonyl)-amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(9,9,10,10,10-pentafluoro-decyl)-amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -[5-(methyl-amino)-pentyl]-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-pyrrolidine-1-yl-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4,4,5,5,5-pentafluoro-pentyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4,4,5,5,6,6,7,7,8,8,9,9,9-tridecafluoro-nonyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(4,4,5,5,6,6,7,7,8,8,9,9,10,10,11,11,11-heptadecafluoro-undecyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

10016449-050706

11 β -{5-[methyl-(3,3,4,4,5,5,6,6,6-nonafluoro-hexyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,8-pentafluoro-octyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{6-[methyl-(7,7,8,8,8-pentafluoro-octyl)-amino]-hexyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,9,9,10,10,10-nonafluoro-decyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(7,7,8,8,9,9,10,10,11,11,12,12,12-tridecafluoro-dodecyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(7,7,8,8,9,9,10,10,11,11,12,12,13,13,14,14-heptadecafluoro-tetradecyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,5-hexafluoro-pent-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,6,6,7,7,8,8,8-dodecafluoro-oct-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3,4,4,5,5,6,6,7,7,8,8,9,9,10,10,10-hexadecafluoro-dec-2-enyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-phenoxy-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[(3-benzyloxy-propyl)-methyl-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

202050 62481001

11 β -{5-[N-methyl-N-3-(4,4,5,5,5-pentafluoropentyloxy)-propylamino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -[9-(4,4,5,5,5-pentafluoropentylsulfinyl)-nonyl]-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(2-p-tolyl-ethyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[2-(4-ethoxy-phenyl)-ethyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-phenyl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-pyridin-3-yl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(3-p-tolyl-propyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[3-(4-chloro-phenyl)-propyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -(5-{[3-(4-ethoxy-phenyl)-propyl]-methyl-amino}-pentyl)-estra-1,3,5(10)-triene-3,17 β -diol

11 β -{5-[methyl-(4-methyl-pentyl)-amino]-pentyl}-estra-1,3,5(10)-triene-3,17 β -diol

7. Use of the compounds of general formula I according to claim 1 for the production of pharmaceutical agents.

8. Pharmaceutical preparations that contain at least one compound of general formula I according to claim 1 as well as a pharmaceutically compatible vehicle.

004849-0000

- (12) INTERNATIONAL APPLICATION PUBLISHED ACCORDING TO THE PATENT
COOPERATION TREATY (PCT)
- (19) World Intellectual Property Organization
International Office
- (43) International publication date
January 4, 2001 (1/4/2001) PCT
- (10) International publication number
WO 01/00652 A2
- (51) International patent classification⁷: C07J
- (21) International file number: PCT/EP00/05969
- (22) International application date: June 26, 2000 (6/26/2000)
- (25) Filing language: German
- (26) Publication language: German
- (30) Priority data: 199 29 715.0 June 24, 1999 (6/24/1999) DE
- (71) Applicant (for all designated countries except for the
U.S.): SCHERING AKTIENGESELLSCHAFT [DE/DE]; Müllerstrasse
178, D-133353 Berlin (DE).
- (72) Inventors; and
- (75) Inventors/applicants (only for the U.S.):
BOHLMANN, Rolf [DE/DE]; Kühler Weg 6A, D-14055 Berlin (DE).
HEINRICH, Nikolaus [DE/DE]; Sponholzstrasse 8, D-12159
Berlin (DE). KROLL, Jorg [DE/DE]; Horst-Kohl-Strasse 3, D-
12157 Berlin (DE). SAUER, Gerhard [DE/DE]; Königsbacher
Zeile 41a, D-13465 Berlin (DE).
- (81) Designated countries (National): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DK, DM, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,

10018429 050700

KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated countries (Regional): ARIPO Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), [continuation on the next page] European Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

-- Without international search report and to be republished after receipt of the report.

To clarify the two-letter code, and the other abbreviations, reference is made to the explanations ("Guidance Notes on Codes and Abbreviations") at the beginning of each regular edition of the PCT Gazette.

(54) Title: 11 β -LONG-CHAIN-SUBSTITUTED ESTRATRIENES, PROCESS FOR THE PRODUCTION, PHARMACEUTICAL PREPARATIONS THAT CONTAIN THESE 11 β -LONG-CHAIN-SUBSTITUTED ESTRATRIENES, AS WELL AS THEIR USE FOR THE PRODUCTION OF PHARMACEUTICAL AGENTS



(I)

(57) **Abstract:** This invention describes the new 11 β -long-chain-substituted estratrienes of general formula (I), in which R¹¹ is a long-chain radical that has a nitrogen atom and optionally a sulfur atom, which in addition can be functionalized in the terminal position with a perfluoroalkyl group or an optionally substituted aryl radical. The compounds have antiestrogenic or tissue-selective estrogenic properties and are suitable for the production of pharmaceutical agents.

10018429 090702

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

11 β LONG-CHAIN SUBSTITUTED ESTRADIENES, METHOD FOR THEIR PRODUCTION, PHARMACEUTICAL PREPARATIONS CONTAINING SAID 11 β LONG-CHAIN SUBSTITUTED ESTRADIENES, AND THEIR USE FOR PRODUCING MEDICAMENTS

the specification of which

☐ is attached hereto

☒ was filed on 26 JUNE 2000 as United States Application Number or PCT International Application Number PCT/EP00/05969 and (if applicable) was amended on _____

I hereby authorize our attorneys to insert the serial number assigned to this application.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 USC §119			
APPLICATION NO.	COUNTRY	DAY/MONTH/YEAR FILED	PRIORITY CLAIMED
199 29 715.0	GERMANY	24 JUNE 1999	YES

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

PROVISIONAL APPLICATION(S) UNDER 35 U.S.C. §119(e)	
APPLICATION NUMBER	FILING DATE

I hereby claim the benefit under 35 U.S.C. §120 of any United States application, or §365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. §112.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

PRIOR U.S./PCT INTERNATIONAL APPLICATION(S) DESIGNATED FOR BENEFIT UNDER 37 U.S.C. §120		
APPLICATION NO.	FILING DATE	STATUS — PATENTED, PENDING, ABANDONED

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith: I. William Millen (19,544); John L. White (17,746); Anthony J. Zelano (27,969); Alan E.J. Branigan (20,565); John R. Moses (24,983); Harry B. Shubin (32,004); Brion P. Heaney (32,542); Richard J. Traverso (30,595); John A. Sopp (33,103); Richard M. Lebovitz (37,067); John H. Thomas (33,460); Catherine M. Joyce (40,668); Nancy J. Axelrod (44,014); James T. Moore (35,619); James E. Ruland (37,432); Jennifer J. Branigan (40,921) and Robert E. McCarthy (46,044)

Send Correspondence to Customer Number 23599



23599

PATENT TRADEMARK OFFICE

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of sole or first inventor (given name, family name)

Rolf BOHLMANN

Signature Rolf Bohlmann

Date 23. Jan '02

Residence
Berlin, Germany

Citizenship
Germany

Post Office Address Kuhler Weg 6A, D-14055 Berlin Germany

Full Name of additional joint inventor (given name, family name)

Nikolaus HEINRICH

Signature Nikolaus Heinrich

Date 23. Jan. 2002

Residence
Berlin, Germany

Citizenship
Germany

Post Office Address Sponholzstrasse 8, D-12159 Berlin Germany

Full Name of additional joint inventor (given name, family name)

Jorg KROLL

Signature Jorg Kroll

Date 23. Jan '2002

Residence
Berlin, Germany

Citizenship
Germany

Post Office Address Horst-Kohl-Strasse 3, D-12157 Berlin Germany

Full Name of additional joint inventor (given name, family name)

Gerhard SAUER

Signature Gerhard Sauer

Date 6. Mar 2002

Residence
Berlin, Germany

Citizenship
Germany

Post Office Address Konigsbacher Zeile 41a, D-13465 Berlin Germany

Full Name of additional joint inventor (given name, family name)

Karl-Heinrich FRITZEMEIER

Signature Karl-Heinrich Fritzemeier

Date 23. Jan '2002

Residence
Berlin, Germany

Citizenship
Germany

Post Office Address Rasbenstrasse 5a, D-13503 Berlin Germany

Full Name of additional joint inventor (given name, family name)

Christa HEGELE-HARTUNG

Signature Christa Hegel-Hartung

Date 29th January, 2002

Residence
Mulheim a.d. Ruhr, Germany

Citizenship
Germany

Post Office Address Wallerback 404, D-45470 Mulheim a.d. Ruhr, Germany

Full Name of additional joint inventor (given name, family name)

Jens HOFFMANN

Signature *Jens Hoffmann*

Date Feb 12, 2002

Residence Mühlenbeck, Germany *Dut*

Citizenship Germany

Post Office Address Walterstrasse 13, D-16567 Mühlenbeck, Germany

Full Name of additional joint inventor (given name, family name)

Rosemarie LICHTNER

Signature *Rosemarie Lichtner*

Date Feb 22, 2002

Residence Berlin, Germany *Dut*

Citizenship Germany

Post Office Address Belzigerstrasse 39, D-10823 Berlin, Germany

Full Name of additional joint inventor (given name, family name)

Ludwig ZORN

Signature *Ludwig Zorn*

Date Feb 18, 2002

Residence Berlin, Germany *Dut*

Citizenship Germany

Post Office Address Osianderweg 45a, D-13509 Berlin, Germany

Full Name of additional joint inventor (given name, family name)

Signature

Date

Residence

Citizenship

Post Office Address

Full Name of additional joint inventor (given name, family name)

Signature

Date

Residence

Citizenship

Post Office Address

Full Name of additional joint inventor (given name, family name)

Signature

Date

Residence

Citizenship

Post Office Address

Additional joint inventors are named on separately numbered sheets attached hereto.